



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Sort

Save

Text

Clip Add

Order

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

PubMed Services

Journal Browser

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: J Invest Surg 1997 Sep-Oct;10(5):295-304

Related Articles, **NEW Books**,
LinkOut**An adult canine model of progressive left ventricular pressure overload.****Keech GB, Smith AC, Swindle MM, Koide M, Carabello BA, DeFreyte G.**

Department of Comparative Medicine, Medical University of South Carolina, Charleston 29425, USA.

This article details the development of a model of progressive left ventricular pressure overload (LVPO) in the adult dog. LVPO was induced by banding the proximal ascending aorta in 69 adult conditioned dogs. The base of the aorta was exposed through a right thoracotomy. A tunnel was created by blunt dissection between the aorta and pulmonary arteries. An aortic band was constructed by passing umbilical tapes through the lumen of gortex tubing. This band was placed through the tunnel, then tied around a balloon dilatation catheter. The distal end of the balloon catheter was closed with an injection cap and positioned in a subcutaneous pocket. Aortic stenosis was induced by filling the balloon catheter with saline. A predetermined amount of LVPO was created by adjusting the amount of aortic stenosis. At 2, 4, and 6 weeks after aortic banding the LVPO was increased by transcutaneous injection of saline into the balloon catheter. At 8 weeks the dogs were evaluated for sufficiently decreased cardiac contractility and used acutely in one of several studies. The article also discusses perioperative management, postoperative care, and complications that were encountered during the development of the model. Postoperative pain was managed by the combined use of preemptive and postoperative opioids, local nerve blocks, and nonsteroidal anti-inflammatory drugs. Notable intraoperative complications included atrial and ventricular arrhythmias and pulmonary artery laceration during the banding procedure. The most significant postoperative complications were aortic ruptures and congestive heart failure. The success rate of this model has increased from 20% (year 1) to 65% (year 3). This success has been attributed to improvements in band design, surgical technique, and postoperative management.